


**Class- X**  
**Mathematics-Basic (241)**  
**Marking Scheme SQP-2020-21**


**Max. Marks: 80**

**Duration:3hrs**

|   |   |                         |
|---|---|-------------------------|
| 1 | $156 = 2^2 \times 3 \times 13$  | 1                       |
| 2 | Quadratic polynomial is given by $x^2 - (a + b)x + ab$<br>$x^2 - 2x - 8$  | 1                       |
| 3 | HCF X LCM = product of two numbers<br>$\text{LCM}(96, 404) = \frac{96 \times 404}{\text{HCF}(96, 404)} = \frac{96 \times 404}{4}$ <p style="text-align: center;">LCM = 9696</p> <p style="text-align: center;"><b>OR</b></p> <p>Every composite number can be expressed (factorized) as a product of primes, and this factorization is unique, apart from the order in which the factors occur.</p> | 1/2<br><br>1/2<br><br>1 |
| 4 | $x - 2y = 0$<br><br>$3x + 4y - 20 = 0$<br><br>$\frac{1}{3} \neq \frac{-2}{4}$<br><br>As, $\frac{a_1}{a_2} \neq \frac{b_1}{b_2}$ is one condition for consistency.<br><br>Therefore, the pair of equations is consistent.  | 1/2<br><br>1/2          |
| 5 | 1   | 1                       |
| 6 | $\theta = 60^\circ$<br>Area of sector = $\frac{\theta}{360^\circ} \pi r^2$<br>$A = \frac{60^\circ}{360^\circ} \times \frac{22}{7} \times (6)^2 \text{ cm}^2$<br>$A = \frac{1}{6} \times \frac{22}{7} \times 36 \text{ cm}^2$<br>$= 18.86 \text{ cm}^2$  | 1/2<br><br>1/2          |

|   |   |   |
|---|---|---|
|   | <b>OR</b>   |   |
|   | <p>Another method-</p> <p>Horse can graze in the field which is a circle of radius 28 cm.</p> <p>So, required perimeter = <math>2\pi r = 2 \cdot \pi(28)</math> cm</p> $= 2 \times \frac{22}{7} \times (28) \text{ cm}$ $= 176 \text{ cm}$  | <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p>   |
| 7 | <p>By converse of Thale's theorem <math>DE \parallel BC</math></p> <p><math>\angle ADE = \angle ABC = 70^\circ</math></p> <p>Given <math>\angle BAC = 50^\circ</math></p> <p><math>\angle ABC + \angle BAC + \angle BCA = 180^\circ</math> (Angle sum prop of triangles)</p> $70^\circ + 50^\circ + \angle BCA = 180^\circ$ $\angle BCA = 180^\circ - 120^\circ = 60^\circ$ <p style="text-align: center;"><b>OR</b></p> <p><math>EC = AC - AE = (7 - 3.5) \text{ cm} = 3.5 \text{ cm}</math></p> $\frac{AD}{BD} = \frac{2}{3} \text{ and } \frac{AE}{EC} = \frac{3.5}{3.5} = 1$ <p>So, <math>\frac{AD}{BD} \neq \frac{AE}{EC}</math></p> <p>Hence, By converse of Thale's Theorem, DE is not Parallel to BC.</p> | <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> |
| 8 | <p>Length of the fence = <math>\frac{\text{Total cost}}{\text{Rate}}</math></p> $= \frac{\text{Rs.5280}}{\text{Rs 24/metre}} = 220 \text{ m}$ <p>So, length of fence = Circumference of the field</p> $\therefore 220\text{m} = 2 \pi r = 2 \times \frac{22}{7} \times r$ <p>So, <math>r = \frac{220 \times 7}{2 \times 22} \text{ m} = 35 \text{ m}</math></p>   | <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p>   |
| 9 |  <p>Sol: <math>\tan 30^\circ = \frac{AB}{BC}</math></p> $1/\sqrt{3} = \frac{AB}{8}$ <p><math>AB = 8 / \sqrt{3} \text{ metres}</math></p> <p>Height from where it is broken is <math>8/\sqrt{3} \text{ metres}</math></p>  | <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p>   |


|    |   |   |
|----|---|---|
| 10 | Perimeter = Area<br>$2\pi r = \pi r^2$<br>$r = 2$ units   | 1   |
| 11 | 3 median = mode + 2 mean  | 1   |
| 12 | 8   | 1   |
| 13 | <p> <math>\frac{a_1}{a_2} \neq \frac{b_1}{b_2}</math> is the condition for the given pair of equations to have unique solution.         </p> <p> <math>\frac{4}{2} \neq \frac{p}{2}</math> </p> <p> <math>p \neq 4</math> </p> <p>           Therefore, for all real values of p except 4, the given pair of equations will have a unique solution.         </p> <p style="text-align: center;"><b>OR</b></p> <p>           Here, <math>\frac{a_1}{a_2} = \frac{2}{4} = \frac{1}{2}</math> </p> <p> <math>\frac{b_1}{b_2} = \frac{3}{6} = \frac{1}{2}</math> and <math>\frac{c_1}{c_2} = \frac{5}{7}</math> </p> <p> <math>\frac{1}{2} = \frac{1}{2} \neq \frac{5}{7}</math> </p> <p> <math>\frac{a_1}{a_2} = \frac{b_1}{b_2} \neq \frac{c_1}{c_2}</math> is the condition for which the given system of equations will represent parallel lines.         </p> <p>           So, the given system of linear equations will represent a pair of parallel lines.         </p> | <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> |
| 14 | <p>           No. of red balls = 3, No. black balls = 5<br/>           Total number of balls = 5 + 3 = 8<br/>           Probability of red balls = <math>\frac{3}{8}</math> </p> <p style="text-align: center;"><b>OR</b></p> <p>           Total no of possible outcomes = 6<br/>           There are 3 Prime numbers, 2,3,5.<br/>           So, Probability of getting a prime number is <math>\frac{3}{6} = \frac{1}{2}</math> </p>  | <p>1/2</p> <p>1/2</p> <p>1/2</p> <p>1/2</p> |

|       |   |   |
|-------|---|---|
| 15    |  <p style="text-align: center;"> <math>\tan 60^\circ = \frac{h}{15}</math><br/> <math>\sqrt{3} = \frac{h}{15}</math><br/> <math>h = 15\sqrt{3} \text{ m}</math> </p>   | <p style="text-align: center;"><math>\frac{1}{2}</math></p> <p style="text-align: center;"><math>\frac{1}{2}</math></p> |
| 16    | 1   | 1   |
| 17 i) | <p>Ans : b)<br/>         Cloth material required = 2X S A of hemispherical dome<br/> <math>= 2 \times 2\pi r^2</math><br/> <math>= 2 \times 2 \times \frac{22}{7} \times (2.5)^2 \text{ m}^2</math><br/> <math>= 78.57 \text{ m}^2</math></p>   | 1   |
| ii)   | a) Volume of a cylindrical pillar = $\pi r^2 h$   | 1   |
| iii)  | <p>b) Lateral surface area = <math>2 \times 2\pi r h</math><br/> <math>= 4 \times \frac{22}{7} \times 1.4 \times 7 \text{ m}^2</math><br/> <math>= 123.2 \text{ m}^2</math></p>   | 1   |
| iv)   | <p>d) Volume of hemisphere = <math>\frac{2}{3} \pi r^3</math><br/> <math>= \frac{2}{3} \times \frac{22}{7} \times (3.5)^3 \text{ m}^3</math><br/> <math>= 89.83 \text{ m}^3</math></p>  | 1   |
| v)    | <p>b)<br/>         Sum of the volumes of two hemispheres of radius 1cm each = <math>2 \times \frac{2}{3} \pi 1^3</math><br/>         Volume of sphere of radius 2cm = <math>\frac{4}{3} \pi 2^3</math><br/>         So, required ratio is <math>\frac{2 \times \frac{2}{3} \pi 1^3}{\frac{4}{3} \pi 2^3} = 1:8</math></p> | <p style="text-align: center;"><math>\frac{1}{2}</math></p> <p style="text-align: center;"><math>\frac{1}{2}</math></p> |

|       |  |                   |
|-------|--|-------------------|
| 18 i) | c) (0,0)   | 1                 |
| ii)   | a) (4,6)   | 1                 |
| iii)  | a) (6,5)   | 1                 |
| iv)   | a) (16,0)  | 1                 |
| v)    | b) (-12,6)   | 1                 |
| 19 i) | c) 90°   | 1                 |
| ii)   | b) SAS   | 1                 |
| iii)  | b) 4 : 9   | 1                 |
| iv)   | d) Converse of Pythagoras theorem  | 1                 |
| v)    | a) 48 cm <sup>2</sup>  | 1                 |
| 20 i) | d) parabola  | 1                 |
| ii)   | a) 2   | 1                 |
| iii)  | b) -1, 3   | 1                 |
| iv)   | c) $x^2 - 2x - 3$  | 1                 |
| v)    | d) 0   | 1                 |
| 21    | <p>Let P(x,y) be the required point. Using section formula</p> $\left\{ \frac{m_1x_2+m_2x_1}{m_1+m_2}, \frac{m_1y_2+m_2y_1}{m_1+m_2} \right\} = (x, y)$ $x = \frac{3(8)+1(4)}{3+1}, \quad y = \frac{3(5)+1(-3)}{3+1}$ $x = 7 \quad y = 3$ <p>(7,3) is the required point</p> | <p>1</p> <p>1</p> |

|    |   |                         |
|----|---|-------------------------|
|    | <b>OR</b>   |                         |
|    | <p>Let P(x, y) be equidistant from the points A(7,1) and B(3,5)</p> <p>Given AP = BP. So, AP<sup>2</sup> = BP<sup>2</sup></p> $(x-7)^2 + (y-1)^2 = (x-3)^2 + (y-5)^2$ $x^2 - 14x + 49 + y^2 - 2y + 1 = x^2 - 6x + 9 + y^2 - 10y + 25$ $x - y = 2$   | 1<br><br>1              |
| 22 | <p>By BPT,</p> $\frac{AM}{MB} = \frac{AL}{LC} \dots\dots\dots(1)$ <p>Also, <math>\frac{AN}{ND} = \frac{AL}{LC} \dots\dots\dots(2)</math></p> <p>By Equating (1) and (2) <math>\frac{AM}{MB} = \frac{AN}{ND}</math></p>  | 1/2<br><br>1/2<br><br>1 |
| 23 | <p>To prove: AB + CD = AD + BC.</p> <div style="text-align: center;"> </div> <p>Proof: AS = AP ( Length of tangents from an external point to a circle are equal)</p> <p>BQ = BP</p> <p>CQ = CR</p> <p>DS = DR</p> <p>AS + BQ + CQ + DS = AP + BP + CR + DR</p> <p>(AS + DS) + (BQ + CQ) = (AP + BP) + (CR + DR)</p> <p>AD + BC = AB + CD</p> | 1<br><br>1              |
| 24 | For the correct construction  | 2                       |



|           |  |   |
|-----------|--|---|
|           | $636 = \frac{n}{2} [ 2x 9 + (n-1) 8]$ $1272 = n [ 18 + 8n -8]$ $1272 = n [10 +8n]$ $8n^2 +10n -1272 =0$ $4n^2 + 5n -636 =0$ $n = \frac{-b \pm \sqrt{b^2 -4ac}}{2a}$ $n = \frac{-5 \pm \sqrt{5^2 -4x 4x(-636)}}{2x4}$ $n = \frac{-5 \pm 101}{8}$ $n = \frac{96}{8} \qquad n = \frac{-106}{8}$ $n = 12 \qquad n = \frac{-53}{4}$ <p>n=12 (since n cannot be negative)</p>  | <p>1/2</p> <p>1/2</p>                   |
| <p>27</p> | <p>Let <math>\sqrt{3}</math> be a rational number.<br/> Then <math>\sqrt{3} = p/q</math> HCF (p,q) =1<br/> Squaring both sides<br/> <math>(\sqrt{3})^2 = (p/q)^2</math><br/> <math>3 = p^2/q^2</math><br/> <math>3q^2 = p^2</math><br/> 3 divides <math>p^2</math> » 3 divides p<br/> 3 is a factor of p<br/> Take <math>p = 3C</math><br/> <math>3q^2 = (3C)^2</math><br/> <math>3q^2 = 9C^2</math><br/> 3 divides <math>q^2</math> » 3 divides q<br/> 3 is a factor of q<br/> Therefore 3 is a common factor of p and q<br/> It is a contradiction to our assumption that p/q is rational.<br/> Hence <math>\sqrt{3}</math> is an irrational number.</p> | <p>1</p> <p>1/2</p> <p>1/2</p> <p>1</p> |
| <p>28</p> |   |   |




|    |  |   |
|----|--|---|
|    | <p>Required to prove :- <math>\angle PTQ = 2\angle OPQ</math></p> <p>Sol :- Let <math>\angle PTQ = \theta</math></p> <p>Now by the theorem <math>TP = TQ</math>. So, <math>\triangle TPQ</math> is an isosceles triangle</p> $\angle TPQ = \angle TQP = \frac{1}{2}(180^\circ - \theta)$ $= 90^\circ - \frac{1}{2}\theta$ $\angle OPT = 90^\circ$ $\angle OPQ = \angle OPT - \angle TPQ = 90^\circ - (90^\circ - \frac{1}{2}\theta)$ $= \frac{1}{2}\theta$ $= \frac{1}{2}\angle PTQ$<br>$\angle PTQ = 2\angle OPQ$ | <p>1</p> <p>1</p> <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> |
| 29 | <p>Let Meena has received x no. of 50 re notes and y no. of 100 re notes. So,</p> $50x + 100y = 2000$ $x + y = 25$ <p>multiply by 50</p> $50x + 100y = 2000$ $50x + 50y = 1250$ <hr style="width: 10%; margin-left: 0;"/> $50y = 750$ $y = 15$<br><p>Putting value of <math>y = 15</math> in equation (2)</p> $x + 15 = 25$ $x = 10$<br><p>Meena has received 10 pieces 50 re notes and 15 pieces of 100 re notes</p>  | <p>1</p> <p>1</p> <p>1</p>  |
| 30 | <p>(i) 10,11,12...90 are two digit numbers. There are 81 numbers. So, Probability of getting a two-digit number = <math>\frac{81}{90} = \frac{9}{10}</math></p> <p>(ii) 1, 4, 9,16,25,36,49,64,81 are perfect squares. So, Probability of getting a perfect square number. = <math>\frac{9}{90} = \frac{1}{10}</math></p> <p>(iii) 5, 10,15....90 are divisible by 5. There are 18 outcomes.. So, Probability of getting a number divisible by 5. = <math>\frac{18}{90} = \frac{1}{5}</math></p>                   | <p>1</p> <p>1</p> <p>1</p>  |



|    |  |   |
|----|--|---|
|    | = $1/(\sec A - \tan A)$ , proved.  |   |
| 33 | <p>Given:-</p> <p>Speed of boat = <math>18 \text{ km/hr}</math><br/> Distance = <math>24 \text{ km}</math></p> <p>Let <math>x</math> be the speed of stream.<br/> Let <math>t_1</math> and <math>t_2</math> be the time for upstream and downstream.<br/> As we know that,</p> <p>speed = distance / time<br/> <math>\Rightarrow</math> time = distance / speed</p> <p>For upstream,<br/> Speed = <math>(18 - x) \text{ km/hr}</math><br/> Distance = <math>24 \text{ km}</math><br/> Time = <math>t_1</math><br/> Therefore,</p> $t_1 = \frac{24}{18 - x}$ <p>For downstream,<br/> Speed = <math>(18 + x) \text{ km/hr}</math><br/> Distance = <math>24 \text{ km}</math><br/> Time = <math>t_2</math><br/> Therefore,</p> $t_2 = \frac{24}{18 + x}$ <p>Now according to the question-</p> $t_1 = t_2 + 1$ $\frac{24}{18 - x} = \frac{24}{18 + x} + 1$ $\Rightarrow \frac{24(18 + x) - 24(18 - x)}{(18 - x)(18 + x)} = 1$ $\Rightarrow 48x = (18 - x)(18 + x)$ $\Rightarrow 48x = 324 + 18x - 18x - x^2$ $\Rightarrow x^2 + 48x - 324 = 0$ $\Rightarrow x^2 + 54x - 6x - 324 = 0$ $\Rightarrow x(x + 54) - 6(x + 54) = 0$ $\Rightarrow (x + 54)(x - 6) = 0$ | <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> |

|  |  |   |
|--|--|---|
|  | <p><math>\Rightarrow x = -54</math> or <math>x = 6</math></p> <p>Since speed cannot be negative.</p> <p><math>\Rightarrow x = -54</math> will be rejected</p> <p><math>\therefore x = 6</math></p> <p>Thus, the speed of stream is <math>6 \text{ km/hr}</math>.</p> <p style="text-align: center;"><b>OR</b></p> <p>Let one of the odd positive integer be <math>x</math><br/> then the other odd positive integer is <math>x+2</math><br/> their sum of squares = <math>x^2 + (x+2)^2</math><br/> <math>= x^2 + x^2 + 4x + 4</math><br/> <math>= 2x^2 + 4x + 4</math></p> <p>Given that their sum of squares = 290<br/> <math>\Rightarrow 2x^2 + 4x + 4 = 290</math><br/> <math>\Rightarrow 2x^2 + 4x = 290 - 4 = 286</math><br/> <math>\Rightarrow 2x^2 + 4x - 286 = 0</math><br/> <math>\Rightarrow 2(x^2 + 2x - 143) = 0</math><br/> <math>\Rightarrow x^2 + 2x - 143 = 0</math><br/> <math>\Rightarrow x^2 + 13x - 11x - 143 = 0</math><br/> <math>\Rightarrow x(x+13) - 11(x+13) = 0</math><br/> <math>\Rightarrow (x-11)(x+13) = 0</math><br/> <math>\Rightarrow (x-11) = 0, (x+13) = 0</math><br/> Therefore, <math>x = 11</math> or <math>-13</math><br/> According to question, <math>x</math> is a positive odd integer.<br/> Hence, We take positive value of <math>x</math><br/> So, <math>x = 11</math> and <math>(x+2) = 11 + 2 = 13</math><br/> Therefore, the odd positive integers are 11 and 13.</p> | <p><math>\frac{1}{2}</math></p> <p>1</p> <p>1</p> <p>1</p> <p>1</p> |
|--|--|---|

34



Let AB and CD be the multi-storeyed building and the building respectively.

Let the height of the multi-storeyed building =  $h$  m and  
the distance between the two buildings =  $x$  m.

$$AE = CD = 8 \text{ m [Given]}$$

$$BE = AB - AE = (h - 8) \text{ m}$$

and

$$AC = DE = x \text{ m [Given]}$$

Also,

$$\angle FBD = \angle BDE = 30^\circ \text{ ( Alternate angles)}$$

$$\angle FBC = \angle BCA = 45^\circ \text{ (Alternate angles)}$$

Now,

In  $\Delta ACB$ ,

$$\Rightarrow \tan 45^\circ = \frac{AB}{AC} \left[ \because \tan \theta = \frac{\text{Perpendicular}}{\text{Base}} \right]$$

$$\Rightarrow 1 = \frac{h}{x}$$

$$\Rightarrow x = h \dots (i)$$

In  $\Delta BDE$ ,

1

 $\frac{1}{2}$ 

1

$$\Rightarrow \tan 30^\circ = \frac{BE}{ED}$$

$$\Rightarrow \frac{1}{\sqrt{3}} = \frac{h-8}{x}$$

$$\Rightarrow x = \sqrt{3}(h-8) \dots \dots \dots (ii)$$

From (i) and (ii), we get,

$$h = \sqrt{3}h - 8\sqrt{3}$$

$$\sqrt{3}h - h = 8\sqrt{3}$$

$$h(\sqrt{3} - 1) = 8\sqrt{3}$$

$$h = \frac{8\sqrt{3}}{\sqrt{3}-1}$$

$$h = \frac{8\sqrt{3}}{\sqrt{3}-1} \times \frac{\sqrt{3}+1}{\sqrt{3}+1}$$


$$h = 4\sqrt{3}(\sqrt{3} + 1)$$

$$h = 12 + 4\sqrt{3} \text{ m}$$

Distance between the two building

$$x = (12 + 4\sqrt{3}) \text{ m} \quad [From(i)]$$

OR



From the figure, the angle of elevation for the first position of the balloon  $\angle EAD = 60^\circ$  and for second position  $\angle BAC = 30^\circ$ . The vertical distance

$$ED = CB = 88.2 - 1.2 = 87 \text{ m.}$$

1

1

1/2

1

|    |  |   |
|----|--|---|
|    | <p>Let AD = x m and AB = y m.</p> <p>Then in right <math>\Delta ADE</math>, <math>\tan 60^\circ = \frac{DE}{AD}</math></p> $\sqrt{3} = \frac{87}{x}$ $x = \frac{87}{\sqrt{3}} \dots\dots\dots(i)$ <p>In right <math>\Delta ABC</math>, <math>\tan 30^\circ = \frac{BC}{AB}</math></p> $\frac{1}{\sqrt{3}} = \frac{87}{y}$ $y = 87\sqrt{3} \dots\dots\dots(ii)$ <p>Subtracting(i) and (ii)</p> $y-x = 87\sqrt{3} - \frac{87}{\sqrt{3}}$ $y-x = \frac{87 \cdot 2 \cdot \sqrt{3}}{\sqrt{3} \cdot \sqrt{3}}$ $y-x = 58\sqrt{3} \text{ m}$ <p>Hence, the distance travelled by the balloon is equal to BD</p> $y-x = 58\sqrt{3} \text{ m.}$ | <p>1</p> <p>1</p> <p>1</p> <p>1</p>   |
| 35 | <p>Let A be the first term and D the common difference of A.P.</p> $T_p = a = A + (p-1)D = (A-D) + pD \quad (1)$ $T_q = b = A + (q-1)D = (A-D) + qD \quad \dots(2)$ $T_r = c = A + (r-1)D = (A-D) + rD \quad \dots(3)$ <p>Here we have got two unknowns A and D which are to be eliminated.</p> <p>We multiply (1),(2) and (3) by <math>q-r, r-p</math> and <math>p-q</math> respectively and add:</p> $a(q-r) = (A-D)(q-r) + Dp(q-r)$ $b(r-p) = (A-D)(r-p) + Dq(r-p)$ $c(p-q) = (A-D)(p-q) + Dr(p-q)$ $a(q-r) + b(r-p) + c(p-q)$ $= (A-D)[q-r+r-p+p-q] + D[p(q-r) + q(r-p) + r(p-q)]$ $= (A-D)(0) + D[pq-pr + qr-pq + rp-rq]$ $= 0$   | <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> <p><math>\frac{1}{2}</math></p> <p>1</p> <p>1</p> |

|  |   |                         |                          |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
|--|---|-------------------------|--------------------------|-------------------------|--------------------------|-----------|---|---------|-----|---------|---------|-------|-------|---------|----|-------|---------|---------|----|---------|--------|---------|---|-------|-----|---------|---|-------|-------|--|--|-----------|--|---|
| 36   | <table border="0"> <tr> <td><b>Height (in cm)</b></td> <td><b>f</b></td> <td><b>C.F.</b></td> </tr> <tr> <td>below 140</td> <td>4</td> <td>4</td> </tr> <tr> <td>140-145</td> <td>7</td> <td>11</td> </tr> <tr> <td>145-150</td> <td>18</td> <td>29</td> </tr> <tr> <td>150-155</td> <td>11</td> <td>40</td> </tr> <tr> <td>155-160</td> <td>6</td> <td>46</td> </tr> <tr> <td>160-165</td> <td>5</td> <td>51</td> </tr> </table>   | <b>Height (in cm)</b>   | <b>f</b>                 | <b>C.F.</b>             | below 140                | 4         | 4 | 140-145 | 7   | 11      | 145-150 | 18    | 29    | 150-155 | 11 | 40    | 155-160 | 6       | 46 | 160-165 | 5      | 51      | 1 |       |     |         |   |       |       |  |  |           |  |   |
|  | <b>Height (in cm)</b>   | <b>f</b>                | <b>C.F.</b>              |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
| below 140  | 4   | 4                       |                          |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
| 140-145  | 7   | 11                      |                          |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
| 145-150  | 18  | 29                      |                          |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
| 150-155  | 11  | 40                      |                          |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
| 155-160  | 6   | 46                      |                          |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
| 160-165  | 5   | 51                      |                          |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
| <p><math>N=51 \Rightarrow</math></p> <p><math>N/2=51/2=25.5</math></p> <p>As 29 is just greater than 25.5, therefore median class is 145-150.</p> $\text{Median} = l + \frac{\left(\frac{N}{2} - C\right)}{f} \times h$ <p>Here, <math>l</math> = lower limit of median class = 145</p> <p><math>C</math> = C.F. of the class preceding the median class = 11</p> <p><math>h</math> = higher limit - lower limit = 150 - 145 = 5</p> <p><math>f</math> = frequency of median class = 18</p> <p><math>\therefore \text{median} =</math></p> $= 145 + \frac{(25.5 - 11)}{18} \times 5$ <p>= 149.03</p> | 1/2   |                         |                          |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
|  | <p>Mean by direct method</p> <table border="0"> <tr> <td><b>Height (in cm)</b></td> <td><b>f</b></td> <td><b><math>x_i</math></b></td> <td><b><math>fx_i</math></b></td> </tr> <tr> <td>below 140</td> <td>4</td> <td>137.5</td> <td>550</td> </tr> <tr> <td>140-145</td> <td>7</td> <td>142.5</td> <td>997.5</td> </tr> <tr> <td>145-150</td> <td>18</td> <td>147.5</td> <td>2655</td> </tr> <tr> <td>150-155</td> <td>11</td> <td>152.5</td> <td>1677.5</td> </tr> <tr> <td>155-160</td> <td>6</td> <td>157.5</td> <td>945</td> </tr> <tr> <td>160-165</td> <td>5</td> <td>162.5</td> <td>812.5</td> </tr> <tr> <td></td> <td></td> <td><math>\sum fx</math></td> <td></td> </tr> </table> $\text{Mean} = \frac{\sum fx}{N}$ <p>= 7637.5/51</p> <p>= 149.75</p> | <b>Height (in cm)</b>   | <b>f</b>                 | <b><math>x_i</math></b> | <b><math>fx_i</math></b> | below 140 | 4 | 137.5   | 550 | 140-145 | 7       | 142.5 | 997.5 | 145-150 | 18 | 147.5 | 2655    | 150-155 | 11 | 152.5   | 1677.5 | 155-160 | 6 | 157.5 | 945 | 160-165 | 5 | 162.5 | 812.5 |  |  | $\sum fx$ |  | 1 |
| <b>Height (in cm)</b>  | <b>f</b>  | <b><math>x_i</math></b> | <b><math>fx_i</math></b> |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
| below 140  | 4   | 137.5                   | 550                      |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
| 140-145  | 7   | 142.5                   | 997.5                    |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
| 145-150  | 18  | 147.5                   | 2655                     |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
| 150-155  | 11  | 152.5                   | 1677.5                   |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
| 155-160  | 6   | 157.5                   | 945                      |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
| 160-165  | 5   | 162.5                   | 812.5                    |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |
|  |   | $\sum fx$               |                          |                         |                          |           |   |         |     |         |         |       |       |         |    |       |         |         |    |         |        |         |   |       |     |         |   |       |       |  |  |           |  |   |